

Origin and characterization of alpha smooth muscle actin-positive cells during murine lung development

Moiseenko A., Kheirollahi V., Chao C., Ahmadvand N., Quantius J., Wilhelm J., Herold S., Ahlbrecht K., Morty R., Rizvanov A., Minoo P., El Agha E., Bellusci S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 The Authors Stem Cells published by Wiley Periodicals, Inc. on behalf of AlphaMed Press
ACTA2 expression identifies pulmonary airway and vascular smooth muscle cells (SMCs) as well as alveolar myofibroblasts (MYF). Mesenchymal progenitors expressing fibroblast growth factor 10 (Fgf10), Wilms tumor 1 (Wt1), or glioma-associated oncogene 1 (Gli1) contribute to SMC formation from early stages of lung development. However, their respective contribution and specificity to the SMC and/or alveolar MYF lineages remain controversial. In addition, the contribution of mesenchymal cells undergoing active WNT signaling remains unknown. Using Fgf10 CreERT2, Wt1 CreERT2, Gli1 CreERT2, and Axin2 CreERT2 inducible driver lines in combination with a tdTomato flox reporter line, the respective differentiation of each pool of labeled progenitor cells along the SMC and alveolar MYF lineages was quantified. The results revealed that while FGF10 + and WT1 + cells show a minor contribution to the SMC lineage, GLI1 + and AXIN2 + cells significantly contribute to both the SMC and alveolar MYF lineages, but with limited specificity. Lineage tracing using the Acta2-CreERT2 transgenic line showed that ACTA2 + cells labeled at embryonic day (E)11.5 do not expand significantly to give rise to new SMCs at E18.5. However, ACTA2 + cells labeled at E15.5 give rise to the majority (85%–97%) of the SMCs in the lung at E18.5 as well as alveolar MYF progenitors in the lung parenchyma. Fluorescence-activated cell sorting-based isolation of different subpopulations of ACTA2 + lineage-traced cells followed by gene arrays, identified transcriptomic signatures for alveolar MYF progenitors versus airway and vascular SMCs at E18.5. Our results establish a new transcriptional landscape for further experiments addressing the function of signaling pathways in the formation of different subpopulations of ACTA2 + cells. Stem Cells 2017;35:1566–1578.

<http://dx.doi.org/10.1002/stem.2615>

Keywords

Alveolar myofibroblast, Lineage tracing, Lung, Lung development, Smooth muscle cell

References

- [1] Kim HY, Pang M-F, Varner VD et al. Localized smooth muscle differentiation is essential for epithelial bifurcation during branching morphogenesis of the mammalian lung. *Dev Cell* 2015;34:719–726.
- [2] Mailleux AA, Kelly R, Veltmaat JM et al. Fgf10 expression identifies parabronchial smooth muscle cell progenitors and is required for their entry into the smooth muscle cell lineage. *Development* 2005;132:2157–2166.

- [3] El Agha E, Al Alam D, Carraro G et al. Characterization of a novel fibroblast growth factor 10 (Fgf10) knock-in mouse line to target mesenchymal progenitors during embryonic development. *PLoS One* 2012;7:e38452.
- [4] El Agha E, Herold S, Al Alam D et al. Fgf10-positive cells represent a progenitor cell population during lung development and postnatally. *Development* 2014;141:296-306.
- [5] Que J, Wilm B, Hasegawa H et al. Mesothelium contributes to vascular smooth muscle and mesenchyme during lung development. *Proc Natl Acad Sci USA* 2008;105:16626-16630.
- [6] Greif DM, Kumar M, Lighthouse JK et al. Radial construction of an arterial wall. *Dev Cell* 2012;23:482-493.
- [7] Dixit R, Ai X, Fine A. Derivation of lung mesenchymal lineages from the fetal mesothelium requires hedgehog signaling for mesothelial cell entry. *Development* 2013;140:4398-4406.
- [8] Grindley JC, Bellusci S, Perkins D et al. Evidence for the involvement of the Gli gene family in embryonic mouse lung development. *Dev Biol* 1997;188:337-348.
- [9] Hynes M, Stone DM, Dowd M et al. Control of cell pattern in the neural tube by the zinc finger transcription factor and oncogene Gli-1. *Neuron* 1997;19:15-26.
- [10] Lee J, Platt KA, Censullo P et al. Gli1 is a target of sonic hedgehog that induces ventral neural tube development. *Development* 1997;124:2537-2552.
- [11] Miller L-AD, Wert SE, Clark JC et al. Role of sonic hedgehog in patterning of tracheal-bronchial cartilage and the peripheral lung. *Dev Dyn* 2004;231:57-71.
- [12] Bai CB, Stephen D, Joyner AL. All mouse ventral spinal cord patterning by Hedgehog is Gli dependent and involves an activator function of Gli3. *Dev Cell* 2004;6:103-115.
- [13] Méthot N, Basler K. An absolute requirement for Cubitus interruptus in Hedgehog signaling. *Development* 2001;128:733-742.
- [14] Moral P-M, del, Langhe SP, De, Sala FG et al. Differential role of FGF9 on epithelium and mesenchyme in mouse embryonic lung. *Dev Biol* 2006;293:77-89.
- [15] Pepicelli CV, Lewis PM, McMahon AP. Sonic hedgehog regulates branching morphogenesis in the mammalian lung. *Curr Biol* 1998;8:1083-1086.
- [16] Li C, Li M, Li S et al. Progenitors of secondary crest myofibroblasts are developmentally committed in early lung mesoderm. *Stem Cells* 2015;33:999-1012.
- [17] Peng T, Tian Y, Boogerd CJ et al. Coordination of heart and lung co-development by a multipotent cardiopulmonary progenitor. *Nature* 2013;advance on.
- [18] Shan L, Subramaniam M, Emanuel RL et al. Centrifugal migration of mesenchymal cells in embryonic lung. *Dev Dyn* 2008;237:750-757.
- [19] Kumar ME, Bogard PE, Espinoza FH et al. Defining a mesenchymal progenitor niche at single-cell resolution. *Science* 2014;346:1258810-1258810.
- [20] Cohen ED, Ihida-Stansbury K, Lu MM et al. Wnt signaling regulates smooth muscle precursor development in the mouse lung via a tenascin C/PDGFR pathway. *J Clin Invest* 2009;119:2538-2549.
- [21] Sheikh AQ, Lighthouse JK, Greif DM. Recapitulation of developing artery muscularization in pulmonary hypertension. *Cell Rep* 2014;6:809-817.
- [22] Boström H, Willetts K, Pekny M et al. PDGF-A signaling is a critical event in lung alveolar myofibroblast development and alveogenesis. *Cell* 1996;85:863-873.
- [23] Lindahl P, Karlsson L, Hellström M et al. Alveogenesis failure in PDGF-A-deficient mice is coupled to lack of distal spreading of alveolar smooth muscle cell progenitors during lung development. *Development* 1997;124:3943-3953.
- [24] McGowan SE, Grossmann RE, Kimani PW et al. Platelet-derived growth factor receptor-alpha-expressing cells localize to the alveolar entry ring and have characteristics of myofibroblasts during pulmonary alveolar septal formation. *Anat Rec (Hoboken)* 2008;291:1649-1661.
- [25] Kelly R, Brown N, Buckingham M. The arterial pole of the mouse heart forms from Fgf10-expressing cells in pharyngeal mesoderm. *Dev Cell* 2001;1:435-440.
- [26] Wendling O, Bornert J-M, Chambon P et al. Efficient temporally-controlled targeted mutagenesis in smooth muscle cells of the adult mouse. *Genesis* 2009;47:14-18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18942088>. Accessed June 7, 2016.
- [27] Volckaert T, Dill E, Campbell A et al. Parabronchial smooth muscle constitutes an airway epithelial stem cell niche in the mouse lung after injury. *J Clin Invest* 2011;121:4409-4419.
- [28] R Development Core Team, 2007. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <http://www.R-project.org>. Accessed June 23, 2015.
- [29] Smyth GK. Limma: Linear models for microarray data. In: *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*. Editors: Gentleman R, Carey VJ, Huber W, Irizarry RA, Dudoit S. New York, NY: Springer-Verlag, 2005:397-420.

- [30] Gentleman RC, Carey VJ, Bates DM et al. Bioconductor: Open software development for computational biology and bioinformatics. *Genome Biol* 2004;5:R80.
- [31] Smyth GK, Speed T. Normalization of cDNA microarray data. *Methods* 2003;31:265-273.
- [32] Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 2004;3:Article3.
- [33] Bellusci S, Furuta Y, Rush MG et al. Involvement of sonic hedgehog (SHH) in mouse embryonic lung growth and morphogenesis. *Development* 1997;124:53-63.
- [34] Dickie R, Wang YT, Butler JP et al. Distribution and quantity of contractile tissue in postnatal development of rat alveolar interstitium. *Anat Rec (Hoboken)* 2008;291:83-93.
- [35] Chen L, Acciani T, Cras T, Le et al. Dynamic regulation of platelet-derived growth factor receptor α expression in alveolar fibroblasts during realveolarization. *Am J Resp Cell Mol Biol* 2012;47:517-527.
- [36] Nakamura E, Nguyen M-T, Mackem S. Kinetics of tamoxifen-regulated Cre activity in mice using a cartilage-specific CreERT to assay temporal activity windows along the proximodistal limb skeleton. *Dev Dyn* 2006;235:2603-2612.
- [37] Danielian PS, Muccino D, Rowitch DH et al. Modification of gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. *Curr Biol* 1998;8:1323-1326.
- [38] Robinson SP, Langan-Fahey SM, Johnson DA et al. Metabolites, pharmacodynamics, and pharmacokinetics of tamoxifen in rats and mice compared to the breast cancer patient. *Drug Metab Disp* 1991;19:36-43.
- [39] Zervas M, Millet S, Ahn S et al. Cell behaviors and genetic lineages of the mesencephalon and rhombomere 1. *Neuron* 2004;43:345-357.
- [40] Ahn S, Joyner AL. Dynamic changes in the response of cells to positive hedgehog signaling during mouse limb patterning. *Cell* 2004;118:505-516.
- [41] Ahn S, Joyner AL. In vivo analysis of quiescent adult neural stem cells responding to sonic hedgehog. *Nature* 2005;437:894-897.
- [42] El Agha E, Moiseenko A, Kheirollahi V et al. Two-way conversion between lipogenic and myogenic fibroblastic phenotypes marks the progression and resolution of lung fibrosis. *Cell Stem Cell* 2017;20:261-273.e3.
- [43] Fatigati V, Murphy RA. Actin and tropomyosin variants in smooth muscles. Dependence on tissue type. *J Biol Chem* 1984;259:14383-14388.
- [44] Owens GK, Kumar MS, Wamhoff BR. Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 2004;84:767-801.
- [45] Du Y, Guo M, Whitsett JA et al. "LungGENS": A web-based tool for mapping single-cell gene expression in the developing lung. *Thorax* 2015;70:1092-1094.
- [46] Li C, Xiao J, Hormi K et al. Wnt5a participates in distal lung morphogenesis. *Dev Biol* 2002;248:68-81. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12142021>. Accessed September 6, 2016.
- [47] Kumawat K, Menzen MH, Bos IST et al. Noncanonical WNT-5A signaling regulates TGF- β -induced extracellular matrix production by airway smooth muscle cells. *FASEB J* 2013;27:1631-1643.
- [48] Ramasamy SK, Mailleux AA, Gupte VV et al. Fgf10 dosage is critical for the amplification of epithelial cell progenitors and for the formation of multiple mesenchymal lineages during lung development. *Dev Biol* 2007;307:237-247.